

CALCULATING r^2

When you calculate a regression line, the value of r^2 may not appear. If not, go to CATALOG and turn DiagnosticsOn.

DISTRIBUTIONS on TI-83

Under DISTR menu (=2nd VARS)

Binomial : $B(n,p)$

Probability distribution	<code>binomialpdf (n,p)</code>
Cumulative distribution	<code>binomialcdf (n,p)</code>
$P(X = x)$	<code>binomialpdf (n,p,x)</code>
$P(X \leq x)$	<code>binomialcdf (n,p,x)</code>

Examples

$$\begin{aligned} \text{binomialpdf}(2, 0.5) &= \{0.25, 0.5, 0.25\} \\ \text{binomialcdf}(2, 0.5) &= \{0.25, 0.75, 1\} \\ \text{binomialpdf}(2, 0.5, 1) &= 0.5 \\ \text{binomialcdf}(2, 0.5, 1) &= 0.75 \end{aligned}$$

Normal : $N(\mu, \sigma)$

Probability density	<code>normalpdf (x, μ, σ)</code>
$P(a \leq X \leq b)$	<code>normalcdf (a, b, μ, σ)</code>

Either σ , or σ and μ , can be omitted, in which case $\mu = 0$ and $\sigma = 1$ are assumed.

Examples

$$\begin{aligned} \text{normalcdf}(-1, 1) &= 0.68 \\ \text{normalcdf}(-2, 2) &= 0.95 \\ \text{normalcdf}(-3, 3) &= 0.997 \end{aligned}$$

Inverse Normal

To find x , given y :

If $P(-\infty < X \leq x) = y$, use `invNorm (y, μ , σ) = x`.

Again, either σ , or μ and σ , can be omitted, in which case $\mu = 0$ and $\sigma = 1$ are assumed.

Examples

$$\begin{aligned} \text{invNorm}(0.5) &= 0 \text{ because } P(-\infty < X \leq 0) = 0.5, \\ \text{invNorm}(0.975) &= 1.96 \approx 2 \text{ because } P(-\infty < X < 2) \approx 0.975 \end{aligned}$$

t -distribution with n Degrees of Freedom: $T(n)$

Probability density	$\text{tpdf}(x, n)$
$P(a \leq X \leq b)$	$\text{tcdf}(a, b, n)$

χ^2 -distribution with n degrees of freedom (df): $\underline{\chi^2}(n)$

Probability density	$\chi^2 \text{ pdf}(x, n)$
$P(a \leq X \leq b)$	$\chi^2 \text{ cdf}(a, b, n)$

F -distribution with n df in numerator and k df in denominator: $F(n, k)$

Probability density	$\text{Fpdf}(x, n, k)$
$P(a \leq X \leq b)$	$\text{Fcdf}(a, b, n, k)$